arXiv:cond-mat/0512193v1 [cond-mat.str-el] 9 Dec 2005

On the Bose-Einstein Condensation of Magnons in Cs_2CuCl_4

S. E. Sebastian[1], V. S. Zapf[2], N. Harrison[2], C. D. Batista[3], P. A. Sharma[2], M. Jaime[2], I. R. Fisher[1], A. Lacerda[2]
[1] *Geballe Laboratory for Advanced Materials and Department of Applied Physics, Stanford University, Stanford, CA 94305*
[2] *MST-NHMFL, Los Alamos National Laboratory, Los Alamos, NM 87545 and*
[3] *Theoretical Division, Los Alamos National Laboratory, Los Alamos, NM 87545*

(Dated: February 2, 2008)

In a recent paper [1], Radu *et al.* report experimental results they claim to support Bose-Einstein condensation (BEC) of magnons in Cs_2CuCl_4. It is true that an experimentally measured critical power law scaling exponent in agreement with the BEC universality class would support the realization of a BEC in magnetic systems that order as a canted antiferromagnet. It can be shown, however, that the claim of Radu *et al.* is overstated in this instance, because their determination of the critical exponent ϕ relies on a model-dependent theoretical approximation to the critical field H_{c1} for which the associated errors are neglected. We show that when these errors are included, the uncertainty in the obtained exponent is so large that the available experimental data cannot be used to differentiate between contending universality classes.

A two parameter fit to only a few data points delineating the critical ordering temperature (T_c) versus magnetic field (H)in the vicinity of the quantum critical point (QCP), to the power law

$$T_c \sim (H - H_{c1})^{\frac{1}{\phi}} \tag{1}$$

with both H_{c1} and the critical exponent ϕ varying has been shown to be unreliable [2, 3]. An independent experimental determination of H_{c1} is therefore required to obtain an accurate estimate of ϕ. Given that neutron scattering measurements on Cs_2CuCl_4 presented in Ref. [4] have provided such a determination, yielding $H_{c1} = 8.44 \pm 0.01$ T, this would be an appropriate value to use in the fit to Eqn. (1). Radu *et al.* instead use a value of 'H_{c1}'$= 8.51$ T in their fit to Eqn. (1), calculated using an approximate theoretical Hamiltonian, that is subsequently assumed to have *zero error* in their analysis. This assumption has two principal inaccuracies. The first is that the model Hamiltonian neglects higher order interactions, thereby introducing an unknown systematic error in 'H_{c1}'. The second is that the exchange couplings used in its computation have significant experimental uncertainty, introducing a large error in 'H_{c1}'. We obtain 'H_{c1}'$= 8.51 \pm 0.12$ T on using the published errors in the exchange interactions [4].

Fig. 1 shows fits of Eqn. (1) to the experimentally measured phase boundary data points using both the experimental value of $H_{c1} = 8.44 \pm 0.01$ T of Coldea *et al.* [4] and the theoretical estimate of 'H_{c1}'$= 8.51 \pm 0.12$ T, yielding $\phi = 2.8 \pm 0.4$ and $\phi = 1.5 \pm 0.9$ respectively, on considering the dominant contribution to the error: $\delta\phi = \frac{d\phi}{dH}|_{H_{c1}} \delta H_{c1}$. The single most important factor



FIG. 1: Points on the ordering phase boundary from the experimental data in [1]. The solid line represents the best fit to ϕ using the experimentally measured value of $H_{c1} = 8.44$ T from [4]. The dashed line represents the best fit to ϕ using the theoretical estimate of 'H_{c1}'$= 8.51$ T as per the analysis technique used in [1]. The inset shows the variation in the fit value of ϕ with the value of H_{c1}.

responsible for the very large error of $\sim$ 60% in the case of the latter as compared to the error of $\sim$ 14% in the former fit, is the extreme sensitivity of the fit ϕ to the theoretical estimate of the critical field 'H_{c1}', as depicted graphically in the inset to Fig. 1.

Given the substantial uncertainty in the value of ϕ that is obtained from a rigorous analysis, it is clear that the available experimental data do not favor the $3d$ BEC universality class ($\phi = 1.5$) over other possibilities, including the $3d$ Ising universality class ($\phi = 2$).

[1] T. Radu *et al.*, Phys. Rev. Lett. **95**, 127202 (2005).
[2] O. Nohadani *et al.*, Phys. Rev. B. **69**, 220402(R) (2004).
[3] S. E. Sebastian *et al.*, Phys. Rev. B. **72**, 100404(R) (2005).
[4] R. Coldea *et al.*, Phys. Rev. Lett. **88**, 137203 (2002).
[5] M. Y. Veillette *et al.*, Phys. Rev. B **71**, 214426 (2005).
[6] S. Sachdev *Quantum Phase Transitions* (Cambridge University Press, 1999).